

Welche Bedeutung haben Oberflächeneigenschaften für das erfolgreiche Tragen von Kontaktlinsen? Teil 1

Literatur

- [1] Nichols JJ, Sinnott LT. Tear film, contact lens, and patient-related factors associated with contact lens-related dry eye. *Invest Ophthalmol Vis Sci*, 2006; 47:1319-28.
- [2] Sintt CW, Longmuir RA. Contact lens strategies for the patient with dry eye. *Ocul Surf*, 2007; 5:294-307.
- [3] Nichols JJ, Mitchell GL, Nichols KK, Chalmers R, Begley C. The performance of the contact lens dry eye questionnaire as a screening survey for contact lens-related dry eye. *Cornea*, 2002; 21:469-75.
 - 1.
- [4] Begley CG, Caffery B, Nichols KK, Chalmers R. Responses of contact lens wearers to a dry eye survey. *Optom Vis Sci*, 2000; 77:40-6.
- [5] Pritchard N, Fonn D, Brazeau D. Discontinuation of contact lens wear: a survey. *Int Contact Lens Clin*, 1999; 26:157-62.
- [6] Rumpakis JMB. New data on contact lens dropouts: an international perspective. *Review of Optometry*, 2010; 147:37-42
- [7] Pritchard N. How can we avoid CL drop-outs? *Optician*, 2001; 5825:222:14-8.
- [8] Young G, Veys J, Pritchard N, Coleman S. A multi-centre study of lapsed contact lens wearers. *Ophthalmic Physiol Opt*, 2002; 22:516-27.
- [9] Ross G, Nasso M, Franklin V, Lydon F, Tighe B. Silicone hydrogels: Trends in products and properties. In: British Contact Lens Association Clinical Conference. Birmingham 2005.
- [10] Roba M, Duncan EG, Hill GA, Spencer ND, Tosatti SGP. Friction Measurements on Contact Lenses in Their Operating Environment. *Tribology Letters*, 2011; 44:387(11).
- [11] Ross G, Tighe B. The extrinsic modification of contact lenses with poly vinyl pyrrolidone and related copolymers. In: British Contact Lens Association Clinical Conference. Birmingham 2010.
- [12] Giles TG. *In vitro* contact angle and coefficient of friction profiles for daily disposable contact lenses. *Acta Ophthalmologica*, 2008; 86(Suppl 243):p. 0.
- [13] Wolffsohn JS, Hunt OA, Chowdhury A. Objective clinical performance of 'comfort-enhanced' daily disposable soft contact lenses. *Cont Lens Anterior Eye*, 2010; 33:88-92.
- [14] Peterson RC, Wolffsohn JS, Nick J, Winterton L, Lally J. Clinical performance of daily disposable soft contact lenses using sustained release technology. *Cont Lens Anterior Eye*, 2006; 29:127-34.
- [15] Koh S, Maeda N, Hamano T, Hirohara Y, Mihashi T, Hori Y, Hosohata J, Fujikado T, Tano Y. Effect of internal lubricating agents of disposable soft contact lenses on higher-order aberrations after blinking. *Eye & Contact Lens*, 2008; 34:100-5.
- [16] Brennan NA. Contact lens-based correlates of soft lens wearing comfort. *Optom Vis Sci*, 2009; 86: E-abstract 90957.
- [17] Berry M, Purslow C, Murphy PJ, Pult H. Contact Lens Materials, Mucin Fragmentation and Relation to Symptoms. *Cornea*, 2012; Publish Ahead of Print: 10.1097/ICO.0b013e3182254009.
- [18] Pult H, Purslow C, Berry M, Murphy PJ. Clinical tests for successful contact lens wear: relationship and predictive potential. *Optom Vis Sci*, 2008; 85:E924-9.

- [19] Pult H, Murphy PJ, Purslow C. A Novel Method to Predict Dry Eye Symptoms in New Contact Lens Wearers. *Optom Vis Sci*, 2009; 86:E1042-50.
- [20] Höh H, Schirra F, Kienecker C, Ruprecht KW. Lid-parallel conjunctival folds are a sure diagnostic sign of dry eye. *Ophthalmologe*, 1995; 92:802-8.
- [21] Pult H, Riede-Pult B. Grading of Lid-Parallel Conjunctival Folds by Novice and Experienced Observers. *Investigative Ophthalmology & Visual Science*, 2011; 52:3739-.
- [22] Pult H, Murphy PJ, Purslow C. Clide-index: a novel method to diagnose and measure contact lens induced dry eye. *Contact Lens and Anterior Eye*, 2010; 33:E-abstract: 256300.
- [23] Korb DR, Herman JP, Greiner JV, Scaffidi RC, Finnemore VM, Exford JM, Blackie CA, Douglass T. Lid wiper epitheliopathy and dry eye symptoms. *Eye Contact Lens*, 2005; 31:2-8.
- [24] Korb DR, Greiner JV, Herman JP, Hebert E, Finnemore VM, Exford JM, Glonek T, Olson MC. Lid-wiper epitheliopathy and dry-eye symptoms in contact lens wearers. *CLAO J*, 2002; 28:211-6.
- [25] Berry M, Pult H, Purslow C, Murphy PJ. Mucins and ocular signs in symptomatic and asymptomatic contact lens wear. *Optom Vis Sci*, 2008; 85:E930-8.
- [26] Pult H, Murphy PJ, Purslow C. The longitudinal impact of soft contact lens wear on lid wiper epitheliopathy and lid-parallel conjunctival folds. In: 6th International Conference on the Tear Film & Ocular Surface: Basic Science and Clinical Relevance. Florence, Italy 2010.
- [27] Korb DR, Herman JP, Solomon JD, Greiner JV, Blackie CA. Lid Wiper Staining and Sequential Fluorescein Instillation. *Invest Ophthalmol Vis Sci*, 2006; 47:ARVO E-Abstract: 242.
- [28] Pult H, Korb DR, Blackie CA, Knop E. About Vital Staining of the Eye and Eyelids. I. The Anatomy, Physiology, and Pathology of the Eyelid Margins and the Lacrimal Puncta by E. Marx. *Optom Vis Sci*, 2010; 87:718-24.
- [29] Korb DR, Blackie CA. Marx's Line of the Upper Lid is Visible in Upgaze Without Lid Eversion. *Eye Contact Lens*, 2010; 36:149-51.
- [30] Rubio EG. Evaluation of upper eye lid inner margin staining after using lubricating eye drops. *Contact Lens and Anterior Eye*, 2011; 34, Supplement 1:S17.
- [31] 2007 report of the international dry eye workshop (DEWS). *Ocul Surf*, 2007; Volume 5.
- [32] Pult H. Dry eye in soft contact lens wearers. *Contact Lens Spectrum*, 2011; 07:26-53.
- [34] Nichols KK, Foulks GN, Bron AJ, Glasgow BJ, Dogru M, Tsubota K, Lemp MA, Sullivan DA. The International Workshop on Meibomian Gland Dysfunction: Executive Summary. *Invest Ophthalmol Vis Sci*, 2011; 52:1922-9.
- [35] Jackson WB. Blepharitis: current strategies for diagnosis and management. *Can J Ophthalmol* 2008; 43:170-9.
- [36] Cohen S. Prospective case history using Systane lubricant eye drops to help reduce symptoms of dry eye associated with CL wear. In: *American Optometric Association*; 2004.
- [37] Andrasko GJ, Ryen KA, Garofalo RJ, Lemp JM. Compatibility of Silicone Hydrogel Lenses With Multi-Purpose Solutions. *Invest Ophthalmol Vis Sci*, 2006;47:ARVO E-Abstract: 2392.
- [38] Carnt N, Jalbert I, Stretton S, Naduvilath T, Papas E. Solution toxicity in soft contact lens daily wear is associated with corneal inflammation. *Optom Vis Sci*, 2007; 84:309-15.
- [39] Garofalo RJ, Dassanayake N, Carey C, Stein J, Stone R, David R. Corneal staining and subjective symptoms with multipurpose solutions as a function of time. *Eye Contact Lens*, 2005; 31:166-74.
- [40] Ward KW. Superficial punctate fluorescein staining of the ocular surface. *Optom Vis Sci* 2008; 85:8-16.

- [41] Morgan PB, Woods CA, Tranoudis IG, Helland M, Efron N, Grupcheva CN, Jones D, Tan K, Pesinova A, Rayn O, Santodomingo J, Malet F, Ve M. International Contact Lens Prescribing in 2011. *CL Spectrum*, 2012; 26-31.
- [42] Dalton K, Subbaraman LN, Rogers R, Jones L. Physical properties of soft contact lens solutions. *Optom Vis Sci*, 2008; 85:122-8.
- [43] Pult H, Murphy PJ, Purslow C. A novel method to predict the dry eye symptoms in new contact lens wearers. *Optom Vis Sci*, 2009; 86:E1042-50.
- [44] Pult H. Endlich wieder Spaß an der Kontaktlinse. DOZ 2010.
- [45] Schiffman RM, Christianson MD, Jacobsen G, Hirsch JD, Reis BL. Reliability and validity of the Ocular Surface Disease Index. *Arch Ophthalmol*, 2000; 118:615-21.