

Optische Kohärenztomographie in der Kontaktlinsenanpassung

Literaturverzeichnis

- [1] Huang D, Swanson EA, Lin CP, et al. Optical coherence tomography. *Science* 1991;254:1178-1181.
- [2] Fercher AF, Hitzenberger CK, Drexler W, Kamp G, Sattmann H. In vivo optical coherence tomography. *Am J Ophthalmol* 1993;116:113-114.
- [3] Scharf R. Optische Kohärenztomographie im Aufwind. 2009.
- [4] Wirbelauer C, Winkler J, Bastian GO, Haberle H, Pham DT. Histopathological correlation of corneal diseases with optical coherence tomography. *Graefes Arch Clin Exp Ophthalmol* 2002;240:727-734.
- [5] Wojtkowski M, Kaluzny B, Zawadzki RJ. New directions in ophthalmic optical coherence tomography. *Optom Vis Sci* 2012;89:524-542.
- [6] Kaluzny BJ, Fojt W, Szkulmowska A, Bajraszewski T, Wojtkowski M, Kowalczyk A. Spectral optical coherence tomography in video-rate and 3D imaging of contact lens wear. *Optom Vis Sci* 2007;84:1104-1109.
- [7] Ortiz S, Siedlecki D, Perez-Merino P, et al. Corneal topography from spectral optical coherence tomography (sOCT). *Biomed Opt Express* 2011;2:3232-3247.
- [8] Siedlecki D, de Castro A, Gamba E, et al. Distortion correction of OCT images of the crystalline lens: gradient index approach. *Optom Vis Sci* 2012;89:E709-718.
- [9] Westphal V, Rollins A, Radhakrishnan S, Izatt J. Correction of geometric and refractive image distortions in optical coherence tomography applying Fermat's principle. *Opt Express* 2002;10:397-404.
- [10] Dunne MC, Davies LN, Wolffsohn JS. Accuracy of cornea and lens biometry using anterior segment optical coherence tomography. *J Biomed Opt* 2007;12:064023.
- [11] Maceo BM, Manss F, Borja D, et al. Contribution of the crystalline lens gradient refractive index to the accommodation amplitude in non-human primates: in vitro studies. *J Vis* 2011;11:23.
- [12] Modis L, Jr., Szalai E, Kolozsvari B, Nemeth G, Vajas A, Berta A. Keratometry evaluations with the Pentacam high resolution in comparison with the automated keratometry and conventional corneal topography. *Cornea* 2012;31:36-41.
- [13] Savini G, Carbonelli M, Sbreglia A, Barboni P, Deluigi G, Hoffer KJ. Comparison of anterior segment measurements by 3 Scheimpflug tomographers and 1 Placido corneal topographer. *J Cataract Refract Surg* 2011;37:1679-1685.
- [14] Stefano VS, Melo Junior LA, Mallmann F, Schor P. Interchangeability between Placido disc and Scheimpflug system: quantitative and qualitative analysis. *Arq Bras Oftalmol* 2010;73:363-366.
- [15] Savini G, Barboni P, Carbonelli M, Hoffer KJ. Agreement between Pentacam and videokeratography in corneal power assessment. *J Refract Surg* 2009;25:534-538.
- [16] Karnowski K, Kaluzny BJ, Szkulmowski M, Gora M, Wojtkowski M. Corneal topography with high-speed swept source OCT in clinical examination. *Biomed Opt Express* 2011;2:2709-2720.
- [17] Szalai E, Berta A, Hassan Z, Modis L, Jr. Reliability and repeatability of swept-source Fourier-domain optical coherence tomography and Scheimpflug imaging in keratoconus. *J Cataract Refract Surg* 2012;38:485-494.
- [18] van der Worp E. Beyond the Cornea Part I. *Global Contact* 2009;42-45.
- [19] Meier D. Das Cornea-Skleral-Profil - Ein Kriterium individueller Kontaktlinsenanpassung. *Die Kontaktlinse* 1992;4-11.
- [20] Bokern S, Hoppe M, Bandlitz S. Genauigkeit und Wiederholbarkeit bei der Klassifizierung des Corneo-Skleral-Profil. *Die Kontaktlinse* 2007;26-28.
- [11] Lathrop KL, Gupta D, Kagemann L, Schuman JS, Sundarraj N. Optical coherence tomography as a rapid, accurate, noncontact method of visualizing the palisades of Vogt. *Invest Ophthalmol Vis Sci* 2012;53:1381-1387.
- [22] Li P, An L, Reif R, Shen TT, Johnstone M, Wang RK. In vivo microstructural and microvascular imaging of the human corneo-scleral limbus using optical coherence tomography. *Biomed Opt Express* 2011;2:3109-3118.
- [23] Bizheva K, Hutchings N, Sorbara L, Moayed AA, Simpson T. In vivo volumetric imaging of the human corneo-scleral limbus with spectral domain OCT. *Biomed Opt Express* 2011;2:1794-1702.
- [24] Chen S, Huang J, Wen D, Chen W, Huang D, Wang Q. Measurement of central corneal thickness by high-resolution Scheimpflug imaging, Fourier-domain optical coherence tomography and ultrasound pachymetry. *Acta Ophthalmol* 2010.
- [25] Huang J, Pesudovs K, Yu A, et al. A comprehensive comparison of central corneal thickness measurement. *Optom Vis Sci* 2011;88:940-949.
- [26] Ge L, Shen M, Tao A, Wang J, Dou G, Lu F. Automatic segmentation of the central epithelium imaged with three optical coherence tomography devices. *Eye Contact Lens* 2012;38:150-157.
- [27] Schmoll T, Unterhuber A, Kolbitsch C, Le T, Stingl A, Leitgeb R. Precise thickness measurements of Bowman's layer, epithelium, and tear film. *Optom Vis Sci* 2012;89:E795-802.
- [28] Wang J, Jiao S, Ruggeri M, Shousha MA, Chen Q. In situ visualization of tears on contact lens using ultra high resolution optical coherence tomography. *Eye Contact Lens* 2009;35:44-49.
- [29] Szczesna-Iskander DH, Iskander DR. Future directions in non-invasive measurements of tear film surface kinetics. *Optom Vis Sci* 2012;89:749-759.
- [30] Ibrahim OM, Dogru M, Takano Y, et al. Application of visante optical coherence tomography tear meniscus height measurement in the diagnosis of dry eye disease. *Ophthalmology* 2010;117:1923-1929.
- [31] Bandlitz S, Purslow C, Murphy PJ, Pult H. Comparison of a new portable digital meniscometer and optical coherence tomography in tear meniscus measurement. 20th Biennial Meeting of the International Society for Eye Research. Berlin, Germany; 2012.
- [32] Bandlitz S, Purslow C, Murphy PJ, Pult H. Evaluation of lower tear meniscus shape with OCT. Congress of the European Association for Vision and Eye Research. Nice, France; 2012.
- [33] Garcia-Lazaro S, Madrid-Costa D, Ferrer-Blasco T, Montes-Mico R, Cervino A. OCT for assessing artificial tears effectiveness in contact lens wearers. *Optom Vis Sci* 2012;89:E62-69.
- [34] Tapaszto B, Veres A, Kosina-Hagyo K, Somfai GM, Nemeth J. OCT Imaging of lid-parallel conjunctival folds in soft contact lens wearers. *Optom Vis Sci* 2011;88:1206-1213.
- [35] Veres A, Tapaszto B, Kosina-Hagyo K, Somfai GM, Nemeth J. Imaging lid-parallel conjunctival folds with OCT and comparing its grading with the slit lamp classification in dry eye patients and normal subjects. *Invest Ophthalmol Vis Sci* 2011;52:2945-2951.
- [36] Hwang HS, Joo C-K. Novel noncontact meibography with anterior segment optical coherence tomography. *Invest Ophthalmol Vis Sci* 2012;53:E-Abstract 590.
- [37] Shen M, Wang MR, Wang J, Yuan Y, Chen F. Entire contact lens imaged in vivo and in vitro with spectral domain optical coherence tomography. *Eye Contact Lens* 2010;36:73-76.
- [38] Shen M, Cui L, Riley C, Wang MR, Wang J. Characterization of soft contact lens edge fitting using ultra-high resolution and ultra-long scan depth optical coherence tomography. *Invest Ophthalmol Vis Sci* 2011;52:4091-4097.