

Dos and Don'ts in der Thematik des Trockenen Auges

Heiko Pult

Literatur

- [1] Stapleton F, Alves M, Bunya VY, et al. TFOS DEWS II Epidemiology Report. *Ocul Surf* 2017;15:334-65.
- [2] Dumbleton K, Caffery B, Dogru M, et al. The TFOS International Workshop on Contact Lens Discomfort: Report of the Subcommittee on Epidemiology. *Invest Ophthalmol Vis Sci* 2013;54:TFOS20-36.
- [3] Pult H, Purslow C, Berry M, et al. Clinical tests for successful contact lens wear: relationship and predictive potential. *Optom Vis Sci* 2008;85:E924-9.
- [4] Schiffman RM, Christianson MD, Jacobsen G, et al. Reliability and validity of the Ocular Surface Disease Index. *Arch Ophthalmol* 2000;118:615-21.
- [5] 2007 report of the international dry eye workshop (DEWS). *Ocul Surf* 2007;Volume 5:65-204.
- [6] Wolffsohn JS, Arita R, Chalmers R, et al. TFOS DEWS II Diagnostic Methodology report. *Ocul Surf* 2017;15:539-74.
- [7] Belmonte C, Nichols JJ, Cox SM, et al. TFOS DEWS II pain and sensation report. *Ocul Surf* 2017;15:404-37.
- [8] Johnson ME, Murphy PJ. Temporal changes in the tear meniscus following a blink. *Experimental Eye Research* 2006;83:517-25.
- [9] Guillon JP. Non-invasive Tearscope Plus routine for contact lens fitting. *Cont Lens Anterior Eye* 1998;21 Suppl 1:S31-40.
- [10] Sickenberger W. Klassifikation von Spaltlampenbefunden: DOZ Verlag; 2014.
- [11] Bandlitz S, Purslow C, Murphy PJ, et al. Evaluation of lower tear meniscus shape with OCT. *Acta Ophthalmologica* 2012;90:249.
- [12] Trunk L, B N, Wolffsohn JS, et al. Evaluation of tear meniscus height using different methods. In: *Contact 17*. Hamburg, Germany; 2017.
- [13] Tian L, Qu JH, Zhang XY, et al. Repeatability and Reproducibility of Noninvasive Keratograph 5M Measurements in Patients with Dry Eye Disease. *J Ophthalmol* 2016;2016:8013621.
- [14] Pult H, Riede-Pult BH. Impact of Conjunctival Folds on Central Tear Meniscus Height. *Investigative Ophthalmology & Visual Science* 2015;56:1459-66.
- [15] Fukuda R, Usui T, Miyai T, et al. Tear meniscus evaluation by anterior segment swept-source optical coherence tomography. *Am J Ophthalmol* 2013;155:620-4 e2.
- [16] Bandlitz S, Purslow C, Murphy PJ, et al. Influence of conjunctival folds on calculated tear meniscus volume along the lower eyelid. *Ocul Surf* 2016;14:377-84.
- [17] Bandlitz S, Purslow C, Murphy PJ, et al. The Relationship between Tear Meniscus Regularity and Conjunctival Folds. *Optom Vis Sci* 2014;91:1037-44.
- [18] Rolando M, Valente C, Barabino S. New test to quantify lipid layer behavior in healthy subjects and patients with keratoconjunctivitis sicca. *Cornea* 2008;27:866-70.
- [19] Pult H, Bandlitz S. Leitfaden Trockenes Auge [Dry eye guideline]. *Der Augenoptiker* 2012;10:71-3.
- [20] Craig JP, Tomlinson A. Importance of the lipid layer in human tear film stability and evaporation. *Optom Vis Sci* 1997;74:8-13.
- [21] Guillon JP. Use of the Tearscope Plus and attachments in the routine examination of the marginal dry eye contact lens patient. *Adv Exp Med Biol* 1998;438:859-67.
- [22] Finis D, Pischel N, Schrader S, et al. Evaluation of lipid layer thickness measurement of the tear film as a diagnostic tool for Meibomian gland dysfunction. *Cornea* 2013;32:1549-53.
- [23] King-Smith PE, Fink BA, Fogt N. Three interferometric methods for measuring the thickness of layers of the tear film. *Optom Vis Sci* 1999;76:19-32.
- [24] McMonnies CW. Tear instability importance, mechanisms, validity and reliability of assessment. *Journal of Optometry* 2018.
- [25] Papas EB, Ciolino JB, Jacobs D, et al. The TFOS International Workshop on Contact Lens Discomfort: Report of the Management and Therapy Subcommittee. *Invest Ophthalmol Vis Sci* 2013;54:TFOS183-203.
- [26] Downie LE. Automated tear film surface quality breakup time as a novel clinical marker for tear hyperosmolarity in dry eye disease. *Investigative Ophthalmology and Visual Science* 2015;56:7260-8.
- [27] Wiedemann D, Sickenberger W, Michel M, et al. Validation of an objective non-invasive method for a qualitative tear film assessment using a corneal topographer. *Academy* 2010.
- [28] Pult H, Riede-Pult B. The new "TF-Scan" application for video keratography: its repeatability and usefulness in tear film break-up time analysis. *Contact Lens and Anterior Eye* 2011;34, Supplement 1:S14.
- [29] Lan W, Lin L, Yang X, et al. Automatic noninvasive tear breakup time (TBUT) and conventional fluorescent TBUT. *Optom Vis Sci* 2014;91:1412-8.
- [30] Best N, Drury L, Wolffsohn JS. Clinical evaluation of the Oculus Keratograph. *Cont Lens Anterior Eye* 2012.
- [31] Pult H, Riede-Pult B. A New Modified Fluorescein Strip: It's Repeatability and Usefulness in Tear Film Break-Up Time Analysis. *Cont Lens Anterior Eye* 2012;35:35- 8.
- [32] Korb DR, Greiner JV, Herman JP, et al. Lid-wiper epitheliopathy and dry-eye symptoms in contact lens wearers. *CLAO J* 2002;28:211-6.
- [33] Varikooty J, Srinivasan S, Jones L. Atypical manifestation of upper lid margin staining in silicone hydrogel lens wearers with symptoms of dry eye. *Contact Lens & Anterior Eye* 2008;31:44-6.

- [34] Korb DR, Herman JP, Solomon JD, et al. Lid Wiper Staining and Sequential Fluorescein Instillation. *Invest Ophthalmol Vis Sci* 2006;47:ARVO E-Abstract: 242.
- [35] Marx E. Über vitale Färbung des Auges und der Augenlider. *Graefe's Archive for Clinical and Experimental Ophthalmology* 1925;116:114-25.
- [36] Delaveris A, Stahl U, Madigan M, et al. Comparative performance of lissamine green stains. *Cont Lens Anterior Eye* 2017.
- [37] Korb DR, Herman JP, Greiner JV, et al. Lid wiper epitheliopathy and dry eye symptoms. *Eye Contact Lens* 2005;31:2-8.
- [38] Marx E. Über vitale Färbungen am Auge und an den Lidern. *Graefe's Archive for Clinical and Experimental Ophthalmology* 1924;114:465-82.
- [39] Yamaguchi M, Kutsuna M, Uno T, et al. Marx line: Fluorescein staining line on the inner lid as indicator of meibomian gland function. *American Journal of Ophthalmology* 2006;141:669-75.
- [40] Bron AJ, Benjamin L, Snibson GR. Meibomian gland disease. Classification and grading of lid changes. *Eye* 1991;5 (Pt 4):395-411.
- [41] Hoh H, Schirra F, Kienecker C, et al. Lid-parallel conjunctival folds are a sure diagnostic sign of dry eye. *Ophthalmologie* 1995; 2:802 - 8.
- [42] Höh H, Schirra F, Kienecker C, et al. Lid-parallel conjunctival folds are a sure diagnostic sign of dry eye. *Ophthalmologie* 1995; 92:802-8.
- [43] Sickenberger W, Pult H, Sickenberger B. LIPCOF and contact lens wearers - A new tool of forecast subjective dryness and degree of comfort of contact lens wearers. *Contactologia* 2000;22:74-9.
- [44] Pult H, Purslow C, Murphy PJ. The relationship between clinical signs and dry eye symptoms. *Eye (Lond)* 2011;25:502-10.
- [45] Pult H. The Repeatability of Lid Parallel Conjunctival Folds and Lid Wiper Epitheliopathy. In: NCC 2018. Veldhoven, NL; 2018.
- [46] Pult H, Riede-Pult B. Grading of Lid-Parallel Conjunctival Folds by Novice and Experienced Observers. *Invest Ophthalmol Vis Sci* 2011;52:3739-.
- [47] Pult H, Bandlitz S. Lid-Parallel Conjunctival Folds and Their Ability to Predict Dry Eye. *Eye Contact Lens* 2017.
- [48] Arita R, Itoh K, Inoue K, et al. Contact Lens Wear Is Associated with Decrease of Meibomian Glands. *Ophthalmology* 2009; 116:379-84.
- [49] Pflugfelder SC, Tseng SC, Sanabria O, et al. Evaluation of subjective assessments and objective diagnostic tests for diagnosing tear-film disorders known to cause ocular irritation. *Cornea* 1998;17:38-56.
- [50] Hom M. Ocular Surface Disease: Does the New Technology Replace the Biomicroscope? *Optom Vis Sci* 2015;82:E-abstract: AS-04.
- [51] Pult H, Riede-Pult B. An Assesment of Subjective and Objective Grading of Meibography Images. *Invest Ophthalmol Vis Sci* 2012; ARVO E-Abstract 53:588.
- [52] Ngo W, Srinivasan S, Schulze M, et al. Repeatability of grading meibomian gland dropout using two infrared systems. *Optom Vis Sci* 2014;91:658-67.
- [53] Pult H, Korb D, Blackie C, et al. Distribution of meibomian gland loss along the nasal, central and temporal segments of the lower eye lid. In: 7th International Conference on the Tear Film & Ocular Surface: Basic Science and Clinical Relevance. Taormina, Sicily, Italy; 2013.
- [54] Pult H. Meibography in clinical practice. *Contact lens & anterior eye : the journal of the British Contact Lens Association* 2013;36:e5.
- [55] Pult H, Riede-Pult BH, Nichols JJ. Relation Between Upper and Lower Lids' Meibomian Gland Morphology, Tear Film, and Dry Eye. *Optom Vis Sci* 2012;89:310-5.
- [56] Pult H, Nichols JJ. A review of meibography. *Optom Vis Sci* 2012;89:760-9.
- [57] Korb DR, Blackie CA. The Number of Meibomian Glands Yielding Liquid Secretion (MGYLS) as a Function of Age and Type of Contact Lens. *Invest Ophthalmol Vis Sci* 2010;51:6248-.
- [58] Blackie CA, Korb DR. Recovery time of an optimally secreting meibomian gland. *Cornea* 2009;28:293-7.
- [59] Murakami D, Blackie CA, Pult H, et al. Meibomian Gland Function Cannot Be Predicted By Meibography In Patients Symptomatic For Dry Eye. *ARVO Meeting Abstracts* 2014;55:27.
- [60] Bilkhu PS, Naroo SA, Wolffsohn JS. Randomised masked clinical trial of the MGRx eyebag for the treatment of meibomian gland dysfunction-related evaporative dry eye. *Br J Ophthalmol* 2014; 98:1707-11.
- [61] Pult H. Irving Fatt Memorial Lecture: Clinical Evaluation of Dry Eye – A Year of Research. In: BCLA Conference. Birmingham; 2014.
- [62] Sullivan BD, Whitmer D, Nichols KK, et al. An objective approach to dry eye disease severity. *Invest Ophthalmol Vis Sci* 2010.
- [63] Karpecki PM. Kanski's Clinical Ophthalmology: A Systematic Approach, 8th ed., Brad Bowling. *Optometry & Vision Science* 2015;92:e386.
- [64] Hom MM, Mastrota KM, Schachter SE. Demodex. *Optom Vis Sci* 2013;90:e198-205.
- [65] Murube J. Demodex hominis. *The Ocular Surface* 2015.
- [66] Jalbert I, Rejab S. Increased numbers of Demodex in contact lens wearers. *Optom Vis Sci* 2015;92:671-8.
- [67] Nicholls SG, Oakley CL, Tan A, et al. Demodex species in human ocular disease: new clinicopathological aspects. *Int Ophthalmol* 2017;37:303-12.
- [68] Zhang XY, Sun XG. [Progress of study on the demodex blepharitis]. *Zhonghua Yan Ke Za Zhi* 2016;52:315-20.
- [69] Lindsley K, Matsumura S, Hatfield E, et al. Interventions for chronic blepharitis. *Cochrane Database Syst Rev* 2012;Cd005556.
- [70] Craig JP, Nichols KK, Akpek EK, et al. TFOS DEWS II Definition and Classification Report. *Ocul Surf* 2017;15:276-83.
- [71] Craig JP, Nelson JD, Azar DT, et al. TFOS DEWS II Report Executive Summary. *Ocul Surf* 2017;15:802-12.
- [72] Tarrant AM, Ryan MF, Hamilton PA, et al. A pictorial review of hypovolaemic shock in adults. *Br J Radiol* 2008;81:252-7.
- [73] Lemp MA. Report of the National Eye Institute/Industry workshop on Clinical Trials in Dry Eyes. *Clao J* 1995;21:221-32.
- [74] Chalmers RL, Begley CG, Caffery B. Validation of the 5-Item Dry Eye Questionnaire (DEQ-5): Discrimination across self-assessed severity and aqueous tear deficient dry eye diagnoses. *Cont Lens Anterior Eye* 2010;33:55-60.
- [75] Pult H, Wolffsohn JS. The Development and Evaluation of the New Ocular Surface Disease Index-6. In: NCC 2018. Veldhoven, NL; 2018.
- [76] Heiligenhaus A, Koch JM, Kruse FE, et al. Diagnosis and differentiation of dry eye disorders. *Ophthalmologie* 1995;92:6-11.
- [77] Jones L, Downie LE, Korb D, et al. TFOS DEWS II Management and Therapy Report. *Ocul Surf* 2017;15:575-628.
- [78] Pult H. News from the TFOS DEWS II Dry Eye Report In: EA00 / 4th Optometry Conference of Central and Southeastern Europe. Pula, Croatia; 2018.