

Vergleichende Messung der nicht invasiven Tränenfilmaufreißzeit (NIBUT) mit vier unterschiedlichen Messgeräten

Quellen

- [1] Willcox MDP, Argueso P, Georgiev GA, et al. TFOS DEWS II Tear Film Report. *Ocul Surf.* 2017;15:366-403.
- [2] Wolffsohn JS, Arita R, Chalmers R, et al. TFOS DEWS II Diagnostic Methodology report. *Ocular Surf.* 2017;15:539-74.
- [3] Wang MT, Murphy PJ, Blades KJ, et al. Comparison of non-invasive tear film stability measurement techniques. *Clin Exp Optom.* 2018;101:13-7.
- [4] Johnson ME, Murphy PJ. The Effect of instilled fluorescein solution volume on the values and repeatability of TBUT measurements. *Cornea.* 2005;24:811-7.
- [5] Mengher LS, Bron AJ, Tonge SR et al. A non-invasive instrument for clinical assessment of the pre-corneal tear film stability. *Current Eye Research.* 1985;4:1-7.
- [6] Mengher LS, Pandher KS, Bron AJ. Non-invasive tearfilm break-up time: Sensitivity and specificity. *Acta Ophthalmologica.* 1986;64:441-4.
- [7] Bron AJ, de Paiva CS, Chauhan SK et al. TFOS DEWS II pathophysiology report. *Ocul Surf.* 2017;15:438-510
- [8] Müller-Treiber A., Mess- und Untersuchungsmethoden in der Kontaktlinsenanpassung, Müller-Treiber, A., Kontaktlinsen-Know-How, Heidelberg, DOZ, 2013;19
- [9] Stapleton F, Alves M, Bunya VY., et al. TFOS DEWS II Epidemiology Report, *Ocul Surf.* 2017;15:334-368
- [10] Jacobi C. Aktuelle Diagnostik und Therapie des Dry Eye Disease. *Concept Ophthalmologie.* 2017;9:10-16
- [11] Craig JP, Nichols KK, Akpek EK, et al. TFOS DEWS II Definition and Classification Report. *Ocul Surf.* 2017;15:276-283.
- [12] Lemp MA, Baudouin C, Baum J., et al. Definition und Klassifikation des trockenen Auges Bericht. *Ocul Surf.* 2007;5:75-92.
- [13] Sheppard AL, Wolffsohn JS. Digital eye strain: prevalence, measurement and amelioration. *BMJ Open Ophthalmology.* 2018; ;3:e000146.
- [14] Smith JA, Albeitz J, Begley C, et al. The Epidemiology of Dry Eye Disease: Report of the Epidemiology Subcommittee of the International Dry Eye WorkShop. *Ocul Surf.* 2007. 93-107.
- [15] Norn MS. Desiccation of the precorneal film. I. Corneal wetting-time. *Acta Ophthalmol (Copenh).* 1969;47:865-80.
- [16] King-Smith PE, Reuter KS, Braun RJ, et al. Tear film breakup and structure studied by simultaneous video recording of fluorescence and tear film lipid layer images. *Invest Ophth Vis Sci.* 2013; 54(7):4900-4909.
- [17] Jandl A, Jaeger K, Bandlitz S. Vergleich verschiedener Messinstrumente zur nicht invasiven Beurteilung der Tränenfilmstabilität (NIBUT). *Die Kontaktlinse.* 2017; 3:6-9.
- [18] Fuhrer C, Vonarburg M. Vergleichsstudie über die non-invasive Beurteilung der Tränenfilmaufreißzeit und Lipidschichtdicke mit dem Keeler Tearscope Plus, dem Polaris und dem Oculus Keratographen. 2016; Bachelorthese (Institut für Optometrie, FHNW Olten)
- [19] Markoulli M, Duong TB, Lin M, et.al. Imaging the Tear Film: A comparison between the subjective Keeler Tearscope Plus and the objective Oculus Keratograph 5M and LipiView Interferometer. *Current Eye Research.* 2018;43:155-162
- [20] Nichols KK, Mitchell GL, Zadnik K. The repeatability of clinical measurements of dry eye. *Cornea.* 2004;23:272-85.
- [21] Fernandez J, Rodriguez-Vallejo M, Martinez J, et al. Agreement and repeatability of objective systems for assessment of the tear film. *Graefes Arch Clin Exp Ophthalmol.* 2018;256:1535-1541.
- [22] Pult H, Riede-Pult B. The new “TF-Scan” application for video keratography: its repeatability and usefulness in tear film break-up time analysis. *Contact Lens and Anterior Eye.* 2011;34e:S14
- [23] Best N, Drury L, Wolffsohn, J.S. Clinical evaluation of the Oculus Keratograph. *Contact Lens and Anterior Eye.* 2012;35:171-174
- [24] Gumus K, Crockett CH, Rao K, et al. Noninvasive assessment of tear stability with the tear stability analysis system in tear dysfunction patients. *Invest Ophthalmol Vis Sci.* 2011;52:456-61.
- [25] Hong J, Sun X, Wei A, et al. Assessment of tear film stability in dry eye with a newly developed keratograph. *Cornea.* 2013;32:716-21.
- [26] Tong L, Teng LS. Review of Literature on Measurements of Non-invasive Break Up Times, Lipid Morphology and Tear Meniscal Height Using Commercially Available Hand-held Instruments. *Curr Eye Res.* 2018;43:567-75.
- [27] Oculus, Gebrauchsanweisung Oculus Keratograph 5M, nd, 19
- [28] Easytear, Instruction Manual Easytear view +; 2015