

Messungen mit verschiedenen Methoden:

Untersuchung tageszeitlicher Schwankungen des Hornhautdurchmessers

Literatur

- [1] Chen Y, Xia X. Comparison of the Orbscan II topographer and the iTrace aberrometer for the measurements of keratometry and corneal diameter in myopic patients. *BMC Ophthalmol.* 2016;16:33.
- [2] Salouti R, Nowroozzadeh MH, Tajbakhsh Z, Bagheri M, Ghoreyshi M, Azizzadeh M, et al. Agreement of Corneal Diameter Measurements Obtained by a Swept-source Biometer and a Scheimpflug-based Topographer. *Cornea.* 2017;36:1373-6.
- [3] Bergmanson JP, Martinez JG. Size does matter: what is the corneo-limbal diameter? *Clin Exp Optom.* 2017;100:522-8.
- [4] Consejo A, Llorens-Quintana C, Radhakrishnan H, Iskander DR. Mean shape of the human limbus. *J Cataract Refract Surg.* 2017;43:667-72.
- [5] Nemeth G, Hassan Z, Szalai E, Berta A, Modis L, Jr. Comparative analysis of white-to-white and angle-to-angle distance measurements with partial coherence interferometry and optical coherence tomography. *J Cataract Refract Surg.* 2010;36:1862-6.
- [6] Pinero DP, Plaza Puche AB, Alio JL. Corneal diameter measurements by corneal topography and angle-to-angle measurements by optical coherence tomography: evaluation of equivalence. *J Cataract Refract Surg.* 2008;34:126-31.
- [7] Bruner C, Skanchy DF, Wooten JP, Chuang AZ, Kim G. Anterior chamber lens sizing: Comparison of white-to-white and scleral spur-to-scleral spur methods. *J Cataract Refract Surg.* 2020;46:95-101.
- [8] Kawamorita T, Uozato H, Kamiya K, Shimizu K. Relationship between ciliary sulcus diameter and anterior chamber diameter and corneal diameter. *J Cataract Refract Surg.* 2010;36:617-24.
- [9] Ghoreishi M, Abdi-Shahshahani M, Peyman A, Pourazizi M. A model for predicting sulcus-to-sulcus diameter in posterior chamber phakic intraocular lens candidates: correlation between ocular biometric parameters. *Int Ophthalmol.* 2019;39:661-6.
- [10] Jose RM, Roy DK. Castroviejo caliper: a useful tool in plastic surgery. *Plast Reconstr Surg.* 2004;114:1006.
- [11] Dominguez-Vicent A, Perez-Vives C, Ferrer-Blasco T, Albaran-Diego C, Montes-Mico R. Interchangeability among five devices that measure anterior eye distances. *Clin Exp Optom.* 2015;98:254-62.
- [12] Baumeister M, Terzi E, Ekici Y, Kohnen T. Comparison of manual and automated methods to determine horizontal corneal diameter. *J Cataract Refract Surg.* 2004;30:374-80.
- [13] Rufer F, Schroder A, Erb C. White-to-white corneal diameter: normal values in healthy humans obtained with the Orbscan II topography system. *Cornea.* 2005;24:259-61.
- [14] Shajari M, Lehmann UC, Kohnen T. Comparison of Corneal Diameter and Anterior Chamber Depth Measurements Using 4 Different Devices. *Cornea.* 2016;35:838-42.
- [15] Venkataraman A, Mardi SK, Pillai S. Comparison of Eye-metrics and Orbscan automated method to determine horizontal corneal diameter. *Indian J Ophthalmol.* 2010;58:219-22.
- [16] Ferrer-Blasco T, Esteve-Taboada JJ, Martinez-Albert N, Alfonso JF, Montes-Mico R. Agreement of white-to-white measurements with the IOLMaster 700, Atlas 9000, and Sirius systems. *Expert Rev Med Devices.* 2018;15:453-9.
- [17] Jesus DA, Kedzia R, Iskander DR. Precise measurement of scleral radius using anterior eye profilometry. *Cont Lens Anterior Eye.* 2017;40:47-52.
- [18] Read SA, Collins MJ. Diurnal variation of corneal shape and thickness. *Optom Vis Sci.* 2009;86:170-80.
- [19] Chakraborty R, Read SA, Collins MJ. Diurnal variations in axial length, choroidal thickness, intraocular pressure, and ocular biometrics. *Invest Ophthalmol Vis Sci.* 2011;52:5121-9.
- [20] Read SA, Collins MJ, Iskander DR. Diurnal variation of axial length, intraocular pressure, and anterior eye biometrics. *Invest Ophthalmol Vis Sci.* 2008;49:2911-8.
- [21] Burfield HJ, Patel NB, Ostrin LA. Ocular Biometric Diurnal Rhythms in Emmetropic and Myopic Adults. *Invest Ophthalmol Vis Sci.* 2018;59:5176-87.
- [22] Salouti R, Nowroozzadeh MH, Zamani M, Ghoreyshi M, Khodaman AR. Comparison of horizontal corneal diameter measurements using the Orbscan IIz and Pentacam HR systems. *Cornea.* 2013;32:1460-4.
- [23] Fernandez J, Rodriguez-Vallejo M, Martinez J, Tauste A, Hueso E, Pinero DP. Confounding sizing in posterior chamber phakic lens selection due to white-to-white measurement bias. *Indian J Ophthalmol.* 2019;67:344-9.
- [24] Ortiz-Toquero S, Zuniga V, Rodriguez G, de Juan V, Martin R. Agreement of corneal measurements between dual rotating Scheimpflug-Placido system and Placido-based topography device in normal and keratoconus eyes. *J Cataract Refract Surg.* 2016;42:1198-206.
- [25] Pinero DP, Molina-Martin A, Camps VJ, de Fez D, Caballero MT. Validation of corneal topographic and aberrometric measurements obtained by color light-emitting diode reflection topography in healthy eyes. *Graefes Arch Clin Exp Ophthalmol.* 2019;257:2437-47.
- [26] du Toit R, Vega JA, Fonn D, Simpson T. Diurnal variation of corneal sensitivity and thickness. *Cornea.* 2003;22:205-9.
- [27] Odenthal MT, Nieuwendaal CP, Venema HW, Oosting J, Kok JH, Kijlstra A. In vivo human corneal hydration control dynamics: a new model. *Invest Ophthalmol Vis Sci.* 1999;40:312-9.
- [28] Kiely PM, Carney LG, Smith G. Diurnal variations of corneal topography and thickness. *Am J Optom Physiol Opt.* 1982;59:976-82.
- [29] Handa T, Mukuno K, Niida T, Uozato H, Tanaka S, Shimizu K. Diurnal variation of human corneal curvature in young adults. *J Refract Surg.* 2002;18:58-62.
- [30] Read SA, Collins MJ, Carney LG. The diurnal variation of corneal topography and aberrations. *Cornea.* 2005;24:678-87.