

Kontaktlinsen-Huckepacksysteme neu betrachtet:

# Einfluss auf die Physiologie des vorderen Auges

## Literatur

- [1] Smelser GK, Ozanics V. Importance of atmospheric oxygen for maintenance of the optical properties of the human cornea. *Science*. 1952;115:140-1.
- [2] Papas EB. The significance of oxygen during contact lens wear. *Cont Lens Anterior Eye*. 2014;37:394-404.
- [3] Dumbleton KA, Chalmers RL, Richter DB, Fonn D. Vascular response to extended wear of hydrogel lenses with high and low oxygen permeability. *Optom Vis Sci*. 2001;78:147-51.
- [4] Holden BA, Mertz GW, McNally JJ. Corneal swelling response to contact lenses worn under extended wear conditions. *Invest Ophthalmol Vis Sci*. 1983;24:218-26.
- [5] Efron N, Morgan PB. Hydrogel contact lens dehydration and oxygen transmissibility. *CLAO J*. 1999;25:148-51.
- [6] DIN EN ISO 18369-4:2018-04.
- [7] Brennan NA. A model of oxygen flux through contact lenses. *Cornea*. 2001;20:104-8.
- [8] Compan V, Aguirre-Arzo M, Edrington TB, Weissman BA. Modeling Corneal Oxygen with Scleral Gas Permeable Lens Wear. *Optom Vis Sci*. 2016;93:1339-48.
- [9] Holden BA, Sweeney DF, Sanderson G. The minimum precorneal oxygen tension to avoid corneal edema. *Invest Ophthalmol Vis Sci*. 1984;25:476-80.
- [10] Harvitt DM, Bonanno JA. Re-evaluation of the oxygen diffusion model for predicting minimum contact lens Dk/t values needed to avoid corneal anoxia. *Optom Vis Sci*. 1999;76:712-9.
- [11] Lira M, Pereira C, Real Oliveira ME, Castanheira EM. Importance of contact lens power and thickness in oxygen transmissibility. *Cont Lens Anterior Eye*. 2015;38:120-6.
- [12] Fatt I. Oxygen-transmissibility considerations for a hard-soft contact-lens combination. *Am J Optom Physiol Opt*. 1977;54:666-72.
- [13] Weissman BA, Ye P. Calculated tear oxygen tension under contact lenses offering resistance in series: piggyback and scleral lenses. *Cont Lens Anterior Eye*. 2006;29:231-7.
- [14] Giasson CJ, Morency J, Melillo M, Michaud L. Oxygen Tension Beneath Scleral Lenses of Different Clearances. *Optom Vis Sci*. 2017;94:466-75.
- [15] Fisher D, Collins MJ, Vincent SJ. Fluid Reservoir Thickness and Corneal Edema during Open-eye Scleral Lens Wear. *Optom Vis Sci*. 2020;97:683-9.
- [16] Michaud L, van der Worp E, Brazeau D, Warde R, Giasson CJ. Predicting estimates of oxygen transmissibility for scleral lenses. *Cont Lens Anterior Eye*. 2012;35:266-71.
- [17] van der Worp E. Editorial: Scleral lenses (are) special. *Contact Lens and Anterior Eye*. 2017;40:271-2.
- [18] Dhallu SK, Huarte ST, Bilkhui PS, Boychev N, Wolffsohn JS. Effect of Scleral Lens Oxygen Permeability on Corneal Physiology. *Optom Vis Sci*. 2020;97:669-75.
- [19] Baldone JA. Piggy-back fitting of contact lenses. *CLAO J*. 1985;11:130-4.
- [20] Polse KA, Decker MR, Sarver MD. Soft and hard contact lenses worn in combination. *Am J Optom Physiol Opt*. 1977;54:660-5.
- [21] Lopez-Alemany A, Gonzalez-Mejome JM, Almeida JB, Parafita MA, Refojo MF. Oxygen transmissibility of piggyback systems with conventional soft and silicone hydrogel contact lenses. *Cornea*. 2006;25:214-9.
- [22] Giasson CJ, Perreault N, Brazeau D. Oxygen tension beneath piggyback contact lenses and clinical outcomes of users. *CLAO J*. 2001;27:144-50.
- [23] Tsubota K, Mashima Y, Murata H, Yamada M. A piggyback contact lens for the correction of irregular astigmatism in keratoconus. *Ophthalmology*. 1994;101:134-9.
- [24] Sengor T, Kurna SA, Aki S, Ozkurt Y. High Dk piggyback contact lens system for contact lens-intolerant keratoconus patients. *Clin Ophthalmol*. 2011;5:331-5.
- [25] Florken LN, Fink BA, Mitchell GL, Hill RM. Corneal oxygen uptake associated with piggyback contact lens systems. *Cornea*. 2007;26:324-35.
- [26] Fleiszig SM, Lee EJ, Wu C, Andika RC, Vallas V, Portoles M, et al. Cytotoxic strains of *Pseudomonas aeruginosa* can damage the intact corneal surface in vitro. *CLAO J*. 1998;24:41-7.
- [27] Robboy MW, Comstock TL, Kalsow CM. Contact lens-associated corneal infiltrates. *Eye Contact Lens*. 2003;29:146-54.
- [28] Lin MC, Polse KA. Hypoxia, overnight wear, and tear stagnation effects on the corneal epithelium: data and proposed model. *Eye Contact Lens*. 2007;33:378-81; discussion 82.
- [29] Kok JH, Boets EP, van Best JA, Kijlstra A. Fluorophotometric assessment of tear turnover under rigid contact lenses. *Cornea*. 1992;11:515-7.
- [30] Paugh JR, Chen E, Heinrich C, Miller H, Gates T, Nguyen AL, et al. Silicone Hydrogel and Rigid Gas-Permeable Scleral Lens Tear Exchange. *Eye Contact Lens*. 2018;44:97-101.
- [31] Vincent SJ, Alonso-Caneiro D, Collins MJ, Beanland A, Lam L, Lim CC, et al. Hypoxic Corneal Changes following Eight Hours of Scleral Contact Lens Wear. *Optom Vis Sci*. 2016;93:293-9.

- [32] Fadel D, Ezekiel DF. Fenestrated Scleral Lenses: Back to the Origins? Review of Their Benefits and Fitting Techniques. *Optom Vis Sci*. 2020;97:807-20.
- [33] Guillon M, Maissa C. Bulbar conjunctival staining in contact lens wearers and non lens wearers and its association with symptomatology. *Cont Lens Anterior Eye*. 2005;28:67-73.
- [34] Graham AD, Truong TN, Lin MC. Conjunctival epithelial flap in continuous contact lens wear. *Optom Vis Sci*. 2009;86:e324-31.
- [35] Sorbara L, Simpson TL, Maram J, Song ES, Bizheva K, Hutchings N. Optical edge effects create conjunctival indentation thickness artefacts. *Ophthalmic Physiol Opt*. 2015;35:283-92.
- [36] Wolffsohn JS, Drew T, Dhallu S, Sheppard A, Hofmann GJ, Prince M. Impact of soft contact lens edge design and mid-peripheral lens shape on the epithelium and its indentation with lens mobility. *Invest Ophthalmol Vis Sci*. 2013;54:6190-7.
- [37] Bandlitz S, Ruland T, Schwarz D, Jndl A, Wolffsohn J, Pult H. Clinical significance of contact lens related changes of ocular surface tissue observed on optical coherence image. NCC. Veldhoven, Netherlands 2018.
- [38] Hall LA, Hunt C, Young G, Wolffsohn J. Factors affecting corneoscleral topography. *Invest Ophthalmol Vis Sci*. 2013;54:3691-701.
- [39] Hall LA, Young G, Wolffsohn JS, Riley C. The influence of corneoscleral topography on soft contact lens fit. *Invest Ophthalmol Vis Sci*. 2011;52:6801-6.
- [40] Bandlitz S, Baumer J, Conrad U, Wolffsohn J. Scleral topography analysed by optical coherence tomography. *Cont Lens Anterior Eye*. 2017.
- [41] Fadel D. The influence of limbal and scleral shape on scleral lens design. *Cont Lens Anterior Eye*. 2018.
- [42] Pinero DP, Martinez-Abad A, Soto-Negro R, Ruiz-Fortes P, Perez-Cambrudi RJ, Ariza-Gracia MA, et al. Differences in corneo-scleral topographic profile between healthy and keratoconus corneas. *Cont Lens Anterior Eye*. 2019;42:75-84.
- [43] Bandlitz S, Esper P, Stein M, Dautzenberg T, Wolffsohn JS. Corneoscleral Topography Measured with Fourier-based Profilometry and Scheimpflug Imaging. *Optom Vis Sci*. 2020.
- [44] Alonso-Caneiro D, Vincent SJ, Collins MJ. Morphological changes in the conjunctiva, episclera and sclera following short-term miniscleral contact lens wear in rigid lens neophytes. *Cont Lens Anterior Eye*. 2016;39:53-61.
- [45] Macedo-de-Araujo RJ, Serramito-Blanco M, van der Worp E, Carracedo G, Gonzalez-Mejome JM. Differences between Inferior and Superior Bulbar Conjunctiva Goblet Cells in Scleral Lens Wearers: A Pilot Study. *Optom Vis Sci*. 2020;97:726-31.