

Presbyopie – eine kurze erfolgreiche Reise zu Kontaktlinsen – Teil 2

Mehr Optionen der Korrektion mit Fortschreiten der Myopie

Literatur

- [16] Woods J, Woods C, Fonn D. Visual Performance of a Multifocal Contact Lens versus Monovision in Established Presbyopes. *Optometry and Vision Science* 2015;92:175–182.
- [19] Lord SR, Dayhew J, Sc BA, Howland A. Multifocal Glasses Impair Edge-Contrast Sensitivity and Depth Perception and Increase the Risk of Falls in Older People. *Journal of the American Geriatrics Society* 2002;50:1760–6.
- [20] Johnson L, Buckley JG, Scally AJ, Elliott DB. Multifocal Spectacles Increase Variability in Toe Clearance and Risk of Tripping in the Elderly. *Invest Ophthalmol Vis Sci* 2007;48:1466–71.
- [21] Evans BJW. Monovision: a review. *Ophthalmic and Physiological Optics* 2007;27:417–39.
- [22] Rajagopalan AS, Bennett ES, Lakshminarayanan V. Visual Performance of Subjects Wearing Presbyopic Contact Lenses. *Optometry and Vision Science* 2006;83:611–615.
- [23] Richdale K, Mitchell GL, Zadnik K. Comparison of Multifocal and Monovision Soft Contact Lens Corrections in Patients With Low-Astigmatic Presbyopia. *Optometry and Vision Science* 2006;83:266–273.
- [24] Situ P, du Toit R, Fonn D, Simpson T. Successful Monovision Contact Lens Wearers Refitted With Bifocal Contact Lenses. *Eye & Contact Lens* 2003;29:181–184.
- [25] Woods J, Woods CA, Fonn D. Early Symptomatic Presbyopes—What Correction Modality Works Best? *Eye & Contact Lens* 2009;35:221–226.
- [26] Winn B, Whitaker D, Elliott DB, Phillips NJ. Factors affecting light-adapted pupil size in normal human subjects. *Invest Ophthalmol Vis Sci* 1994;35:1132–7.
- [27] Cakmak HB, Cagil N, Simavli H, Duzen B, Simsek S. Refractive Error May Influence Mesopic Pupil Size. *Current Eye Research* 2010;35:939–939.
- [28] Guillon M, Dumbleton K, Theodoratos P, Gobbe M, Wooley CB, Moody K. The Effects of Age, Refractive Status, and Luminance on Pupil Size. *Optom Vis Sci* 2016;93:1093–100.
- [29] Papadatou E, Del Águila-Carrasco AJ, Esteve-Taboada JJ, Madrid-Costa D, Cerviño-Expósito A. Objective assessment of the effect of pupil size upon the power distribution of multifocal contact lenses. *Int J Ophthalmol* 2017;10:103–8.
- [30] Fedtke C, Ehrmann K, Thomas V, Bakaraju RC. Association between multifocal soft contact lens decentration and visual performance. *Clin Optom (Auckl)* 2016;8:57–69.
- [31] Stapleton F, Edwards K, Keay L, Naduvilath T, Dart JKG, Brian G, Holden B. Risk Factors for Moderate and Severe Microbial Keratitis in Daily Wear Contact Lens Users. *Ophthalmology* 2012;119:1516–21.
- [32] Efron N, Brennan NA, Chalmers RL, Jones L, Lau C, Morgan PB, Nichols JJ, Szczotka-Flynn LB, Willcox MD. Thirty years of ‘quiet eye’ with etafilcon A contact lenses. *Contact Lens and Anterior Eye* 2020;43:285–97.
- [33] Chalmers RL, Hickson-Curran SB, Keay L, Gleason WJ, Albright R. Rates of Adverse Events With Hydrogel and Silicone Hydrogel Daily Disposable Lenses in a Large Postmarket Surveillance Registry: The TEMPO Registry. *Invest Ophthalmol Vis Sci* 2015;56:654–63.
- [34] Steiner TF. Contact Lens Patients Are More Valuable than Eyeglasses-Only Patients. *Review of Optometric Business* November 2016:4.