

## Literatur

- [1] Heiligenhaus A et al. Diagnosis and differentiation of dry eye disorders. *Ophthalmologe*, 1995. 92(1): 6-11.
- [2] 2007 report of the international dry eye workshop (DEWS). *Ocul Surf*, 2007, Vol. 5.
- [3] Tiffany JM. The normal tear film. *Dev Ophthalmol*, 2008. 41: 1-20.
- [4] Craig JP, Tomlinson A. Importance of the lipid layer in human tear film stability and evaporation. *Optom Vis Sci*, 1997. 74(1): 8-13.
- [5] Kimball SH, King-Smith PE, Nichols JJ. Evaporation is the Primary Mechanism of Tear Film Thinning between Blinks. *Invest Ophthalmol Vis Sci*, 2010.
- [6] Bischoff G, Khaireddin R. Lipidsubstitution bei kontaktlinsenassoziiertem Trockenen Auge. *Aktuelle Kontaktologie*, 2011. 7: 24-28.
- [7] McCulley JP, Shine WE. The lipid layer: the outer surface of the ocular surface tear film. *Biosci Rep*, 2001. 21(4): 407-18.
- [8] Greiner JV et al. Meibomian gland phospholipids. *Curr Eye Res*, 1996. 15(4): 371-5.
- [9] Khaireddin R, Schmidt KG. Vergleichende Untersuchung zur Therapie des evaporativen trockenen Auges. *Klin. Monatsbl Augenheilkd*, 2010. 227(2): 128-34.
- [10] Craig JP et al. Effect of a liposomal spray on the pre-ocular tear film. *Cont Lens Anterior Eye*, 2010. 33(2): 83-7.
- [11] Dausch D et al. Comparative study of treatment of the dry eye syndrome due to disturbances of the tear film lipid layer with lipid-containing tear substitutes. *Klin Monatsbl Augenheilkd*, 2006. 223(12): 974-83.
- [12] Lee S et al. A New Therapy Concept with a Liposome Eye Spray for the Treatment of the Dry Eye. *Klin Monatsbl Augenheilkd*, 2004. 221: 825-836.
- [13] Schiffman RM et al. Reliability and validity of the Ocular Surface Disease Index. *Arch Ophthalmol*, 2000. 118(5): 615-21.
- [14] The Declaration of Helsinki. *World Med J*, 1982. 29(6): 86-8.
- [15] Pult H, Riede-Pult B. The new 'TF-Scan' application for video keratography: its repeatability and usefulness in tear film break-up time analysis in BCLA Conference 2011, Manchester.
- [16] Schirra F et al. Using LIPCOF (lid-parallel conjunctival fold) for assessing the degree of dry eye, it is essential to observe the exact position of that specific fold. *Adv Exp Med Biol*, 1998. 438: 853-8.
- [17] Höh H et al. Lid-parallel conjunctival folds are a sure diagnostic sign of dry eye. *Ophthalmologe*, 1995. 92(6): 802-8.
- [18] Pult H et al. Clinical tests for successful contact lens wear: relationship and predictive potential. *Optom Vis Sci*, 2008. 85(10): E924-9.

[19] Pult H, Murphy PJ, Purslow C. A Novel Method to Predict Dry Eye Symptoms in New Contact Lens Wearers. *Optom Vis Sci*, 2009. 86(9): E1042-1050.

[20] Berry M et al. Mucins and ocular signs in symptomatic and asymptomatic contact lens wear. *Optom Vis Sci*, 2008. 85(10): E930-8.

[21] Santodomingo-Rubido J, Wolffsohn J, Gilmartin B. Conjunctival epithelial flaps with 18 months of silicone hydrogel contact lens wear. *Eye Contact Lens*, 2008. 34(1): 35-8.

[22] Markoulli M et al. Resolution and Clinical Characteristics of Conjunctival "Flaps". *Invest Ophthalmol Vis Sci*, 2007. 48(5): ARVO E-Abstract: 5391.

[23] Thota S et al. Conjunctival Flaps in Silicone Hydrogel Lens Wearers. *Invest Ophthalmol Vis Sci*, 2006. 47(5): ARVO E-Abstract: 82.

[24] Meng CL. Conjunctival Epithelial Flaps: What Are They and Do We Need to Worry? 05/2006.

[25] Pult H, Riede-Pult B. Grading of Lid-Parallel Conjunctival Folds by Novice and Experienced Invest Ophthalmol Vis Sci, 2011. 52(6): ARVO E-Abstract: 3739.

[26] Pult H, Purslow C, Murphy PJ. The relationship between clinical signs and dry eye symptoms. *Eye (Lond)*, 2011. 25(4): 502-10.

[27] Grein HJ et al. Measuring contrast sensitivity of the human eye with modified Siemens stars. *Optom Vis Sci*, 2010. 87(7): 501-12.

[28] Arnold S et al. Simultaneous examination of tear film break-up and the lipid layer of the human eye: a novel sensor design (Part 1). *Z Med Phys*, 2010. 20(4): 309-15.

[29] Arnold S, Bruenner H, Langenbucher A. Simultaneous examination of tear film break-up and the lipid layer of the human eye: a novel model eye for time course simulation of physiologic tear film behavior (Part 2). *Z Med Phys*, 2010. 20(4): 316-9.

[30] Bende T et al. Contact Free Break Up Time (BUT) Measurement Using a Video Topography System. *Invest Ophthalmol Vis Sci*, 2009. 50(5): ARVO E-Abstract: 524.

[31] Goto T et al. Tear film stability analysis system: introducing a new application for videokeratography. *Cornea*, 2004. 23(8 Suppl): 65-70.

[32] Elliott M et al. Analysis of the repeatability of tear break-up time measurement techniques on asymptomatic subjects before, during and after contact lens wear. *Contact Lens and Anterior Eye*, 1998. 21(4): 98-103.

[33] Wiedemann D et al. Validation of an objective non-invasive method for a qualitative tear film assessment using a corneal topographer. *Academy*, 2010.

[34] Wiedemann D et al. Development and realization of a non-invasive method for tear film assessment by using a corneal topographer. *Academy* 2010.

[35] Pult H, Gill F, Riede-Pult BH. Effect of Three Different Liposomal Eye Sprays on Ocular Comfort and Tear Film, *Cont Lens Anterior Eye*. 2012. (7) e-pub ahead of print].

- [36] Singh O. et al. Chamomile (*Matricaria chamomilla L.*): An overview. *Pharmacogn Rev*, 2011. 5(9): 82-95.
- [37] Paulsen E, Otkjaer A, Andersen KE. The coumarin herniarin as a sensitizer in German chamomile [*Chamomilla recutita (L.) Rauschert, Compositae*]. *Contact Dermatitis*, 2010. 62(6): 338-42.
- [38] Vogel R et al. Demonstration of efficacy in the treatment of dry eye disease with 0.18% sodium hyaluronate ophthalmic solution (vismed, rejena). *Am J Ophthalmol*, 2010. 149(4): 594-601.
- [39] Johnson ME, Murphy PJ, Boulton M. Effectiveness of sodium hyaluronate eyedrops in the treatment of dry eye. *Graefes Arch Clin Exp Ophthalmol*, 2006. 244(1): 109-12.
- [40] O'Brien PD, Collum LM. Dry eye: diagnosis and current treatment strategies. *Curr Allergy Asthma Rep*, 2004. 4(4): 314-9.
- [41] Nichols KK et al. The International Workshop on Meibomian Gland Dysfunction: Executive Summary. *Invest Ophthalmol Vis Sci*, 2011. 52(4): 1922-1929.
- [42] Asari A et al. Hyaluronan, CD44 and fibronectin in rabbit corneal epithelial wound healing. *Japanese Journal of Ophthalmology*, 1996. 40(1): 18-25.
- [43] Sickenberger W, Pult H, B. Sickenberger B. LIPCOF and contact lens wearers - A new tool of forecast subjective dryness and degree of comfort of contact lens wearers. *Contactologia*, 2000. 22(2): 74-79.
- [44] Goto E et al. Impaired functional visual acuity of dry eye patients. *Am J Ophthalmol*, 2002. 133(2): 181-6.
- [45] McMonnies CW. Key questions in a dry eye history. *J Am Optom Assoc*, 1986. 57(7): 512-7.
- [46] McMonnies CW, Ho A. Responses to a dry eye questionnaire from a normal population. *J Am Optom Assoc*, 1987. 58(7): 588-91.
- [47] Wang HF, Fukuda M, Shimomura Y. Diagnosis of dry eye. *Semin Ophthalmol*, 2005. 20(2): 53-62.
- [48] Guillon MPFF et al. Preocular Tear Film Characteristics of Nonwearers and Soft Contact Lens Wearers. *Optom Vis Sci*, 1997. 74(5): 273-279.
- [49] Johnson ME, Murphy PJ. The Effect of instilled fluorescein solution volume on the values and repeatability of TBUT measurements. *Cornea*, 2005. 24(7): 811-7.