

# Hornhautsensibilität

## Teil 3: Das trockene Auge, Keratokonus, refraktive Chirurgie, Keratoplastik und systemische Erkrankungen

Daniela S. Nosch

### Literatur

- [1] Lemp MA, Baudouin MCC, MD, et al. The definition and classification of dry eye disease: report of the Definition and Classification Subcommittee of the International Dry Eye WorkShop (2007). *Ocul Surf*. 2007;5(2):75–92.
- [2] Stern ME, Gao J, Siemasko KF, Beuerman RW, Pflugfelder SC. The role of the lacrimal functional unit in the pathophysiology of dry eye. *Exp Eye Res*. 2004;78(3):409–416. doi:10.1016/j.exer.2003.09.003.
- [3] Dartt DA. Neural regulation of lacrimal gland secretory processes: relevance in dry eye diseases. *Prog Retin Eye Res*. 2009;28(3):155–177. doi:10.1016/j.preteyeres.2009.04.003.
- [4] Jordan A, Baum J. Basic tear flow. Does it exist? *Ophthalmology*. 1980;87(9):920–930.
- [5] Mathers WD. Why the eye becomes dry: a cornea and lacrimal gland feedback model. *CLAO J*. 2000;26(3):159–165.
- [6] Stern ME, BEUERMAN RW, Fox RI, Gao J, Mircheff AK, Pflugfelder SC. A unified theory of the role of the ocular surface in dry eye. *Adv Exp Med Biol*. 1998;438:643–651.
- [7] Acosta MC, Peral A, Luna C, Pintor J, Belmonte C, Gallar J. Tear secretion induced by selective stimulation of corneal and conjunctival sensory nerve fibers. *Invest Ophthalmol Vis Sci*. 2004;45(7):2333–2336.
- [8] Adatia FA, Michaeli-Cohen A, Naor J, Caffery B, Bookman A, Slomovic A. Correlation between corneal sensitivity, subjective dry eye symptoms and corneal staining in Sjögren's syndrome. *Can J Ophthalmol*. 2004;39(7):767–771.
- [9] Xu KP, Yagi Y, Tsubota K. Decrease in corneal sensitivity and change in tear function in dry eye. *Cornea*. 1996;15(3):235–239.
- [10] Hoşal BM, Ornek N, Zilelioğlu G, Elhan AH. Morphology of corneal nerves and corneal sensation in dry eye: a preliminary study. *Eye (Lond)*. 2005;19(12):1276–1279. doi:10.1038/sj.eye.6701760.
- [11] Labbé A, Alalwani H, Van Went C, Brasnu E, Georgescu D, Baudouin C. The relationship between subbasal nerve morphology and corneal sensation in ocular surface disease. *Invest Ophthalmol Vis Sci*. 2012;53(8):4926–4931. doi:10.1167/iovs.11-8708.
- [12] Nishida T, Chikama T-I, Sawa M, Miyata K, Matsui T, Shigeta K. Differential contributions of impaired corneal sensitivity and reduced tear secretion to corneal epithelial disorders. *Jpn J Ophthalmol*. 2012;56(1):20–25. doi:10.1007/s10384-011-0105-4.
- [13] Bourcier T, Acosta MC, Borderie V, et al. Decreased corneal sensitivity in patients with dry eye. *Invest Ophthalmol Vis Sci*. 2005;46(7):2341–2345. doi:10.1167/iovs.04-1426.
- [14] Benítez-Del-Castillo JM, Acosta MC, Wassfi MA, et al. Relation between corneal innervation with confocal microscopy and corneal sensitivity with noncontact esthesiometry in patients with dry eye. *Invest Ophthalmol Vis Sci*. 2007;48(1):173–181. doi:10.1167/iovs.06-0127.
- [15] Benitez-del-Castillo JM, del Rio T, Iradier T, Hernández JL, Castillo A, Garcia-Sanchez J. Decrease in tear secretion and corneal sensitivity after laser in situ keratomileusis. *Cornea*. 2001;20(1):30–32.
- [16] Situ P, Simpson TL, Jones LW, Fonn D. Conjunctival and corneal hyperesthesia in subjects with dryness symptoms. *Optom Vis Sci*. 2008;85(9):867–872. doi:10.1097/OPX.0b013e3181852788.
- [17] De Paiva CS, Pflugfelder SC. Corneal epitheliopathy of dry eye induces hyperesthesia to mechanical air jet stimulation. *Am J Ophthalmol*. 2004;137(1):109–115.
- [18] Tuisku IS, Kontinen YT, Kontinen LM, Tervo TM. Alterations in corneal sensitivity and nerve morphology in patients with primary Sjögren's syndrome. *Exp Eye Res*. 2008;86(6):879–885. doi:10.1016/j.exer.2008.03.002.
- [19] Situ P, Simpson TL, Fonn D, Jones LW. Conjunctival and corneal pneumatic sensitivity is associated with signs and symptoms of ocular dryness. *Invest Ophthalmol Vis Sci*. 2008;49(7):2971–2976. doi:10.1167/iovs.08-1734.
- [20] Golebiowski B, Chim K, So J, Jalbert I. Lid Margins: Sensitivity, Staining, Meibomian Gland Dysfunction, and Symptoms. *Optom Vis Sci*. 2012. doi:10.1097/OPX.0b013e3182693cef.
- [21] Belmonte C, Acosta MC, Gallar J. Neural basis of sensation in intact and injured corneas. *Exp Eye Res*. 2004;78(3):513–525.
- [22] Gallar J, Pozo MA, Tuckett RP, Belmonte C. Response of sensory units with unmyelinated fibres to mechanical, thermal and chemical stimulation of the cat's cornea. *J Physiol (Lond)*. 1993;468:609–622.
- [23] Murphy PJ, Patel S, Morgan PB, Marshall J. The minimum stimulus energy required to produce a cooling sensation in the human cornea. *Ophthalmic Physiol Opt*. 2001;21(5):407–410.
- [24] Acosta MC, Belmonte C, Gallar J. Sensory experiences in humans and single-unit activity in cats evoked by polymodal stimulation of the cornea. *J Physiol (Lond)*. 2001;534(Pt. 2):511–525.
- [25] Belmonte C, Garcia-Hirschfeld J, Gallar J. Neurobiology of Ocular Pain. *Prog Retin Eye Res*. 1997;16(1):117–156.
- [26] Belmonte C, Aracil A, Acosta M, Luna C, Gallar J. Nerves and sensations from the eye surface. *Ocul Surf*. 2004;2(4):248–253.
- [27] Millodot M, Owens H. Sensitivity and fragility in keratoconus. *Acta Ophthalmologica*. 1983;61(5):908–917.
- [28] Simo Mannion L, Tromans C, O'Donnell C. An evaluation of corneal nerve morphology and function in moderate keratoconus. *Cont Lens Anterior Eye*. 2005;28(4):185–192. doi:10.1016/j.clae.2005.10.005.
- [29] Patel DV, Ku JYF, Johnson R, McGhee CNJ. Laser scanning in vivo confocal microscopy and quantitative aesthesiometry reveal decreased corneal innervation and sensation in keratoconus. *Eye*. 2009;23(3):586–592. doi:10.1038/eye.2008.52.
- [30] Krachmer JH, Feder RS, Belin MW. Keratoconus and related noninflammatory corneal thinning disorders. *Surv Ophthalmol*. 1984;28(4):293–322.
- [31] Patel DV, McGhee CNJ. Mapping the corneal sub-basal nerve plexus in keratoconus by in vivo laser scanning confocal microscopy. *Invest Ophthalmol Vis Sci*. 2006;47(4):1348–1351. doi:10.1167/iovs.05-1217.
- [32] Cho KJ, Mok JW, Choi MY, Kim J-Y, Joo C-K. Changes in corneal sensation and ocular surface in patients with asymmetrical keratoconus. *Cornea*. 2013;32(2):205–210. doi:10.1097/ICO.0b013e3182632c07.
- [33] Xia Y, Chai X, Zhou C, Ren Q. Corneal nerve morphology and sensitivity changes after ultraviolet A/riboflavin treatment. *Exp Eye Res*. 2011;93(4):541–547. doi:10.1016/j.exer.2011.06.021.
- [34] Kontadakis GA, Kymionis GD, Kankariya VP, Pallikaris AI. Effect of Corneal Collagen Cross-Linking on Corneal Innervation, Corneal Sensitivity, and Tear Function of Patients with Keratoconus. *Ophthalmology*. 2013. doi:10.1016/j.optha.2012.10.012.

- [35] Petznick, Chew, Hall, et al. Comparison of corneal sensitivity, tear function and corneal staining following laser in situ keratomileusis with two femtosecond laser platforms. *OPHTH*. 2013;591. doi:10.2147/OPHTH.S42266.
- [36] Li M, Zhao J, Shen Y, et al. Comparison of Dry Eye and Corneal Sensitivity between Small Incision Lenticule Extraction and Femtosecond LASIK for Myopia. *PLoS ONE*. 2013;8(10):e77797. doi:10.1371/journal.pone.0077797.
- [37] Darwish T, Brahma A, O'Donnell C, Efron N. Subbasal nerve fiber regeneration after LASIK and LASEK assessed by noncontact esthesiometry and in vivo confocal microscopy: prospective study. *J Cataract Refract Surg*. 2007;33(9):1515–1521. doi:10.1016/j.jcrs.2007.05.023.
- [38] Darwish T, Brahma A, Efron N, O'Donnell C. Subbasal nerve regeneration after LASEK measured by confocal microscopy. *Journal of refractive surgery (Thorofare, NJ : 1995)*. 2007;23(7):709–715.
- [39] Bragheeth MA, Dua HS. Corneal sensation after myopic and hyperopic LASIK: clinical and confocal microscopic study. *British Journal of Ophthalmology*. 2005;89(5):580–585. doi:10.1136/bjo.2004.046888.
- [40] Gallar J, Acosta MC, Moilanen JAO, Holopainen JM, Belmonte C, Tervo TMT. Recovery of corneal sensitivity to mechanical and chemical stimulation after laser in situ keratomileusis. *Journal of refractive surgery (Thorofare, NJ : 1995)*. 2004;20(3):229–235.
- [41] Pérez-Santonja JJ, Sakla HF, Cardona C, Chipont E, Alió JL. Corneal sensitivity after photorefractive keratectomy and laser in situ keratomileusis for low myopia. *Am J Ophthalmol*. 1999;127(5):497–504.
- [42] Murphy PJ, Corbett MC, O'Brart DP, Verma S, Patel S, Marshall J. Loss and recovery of corneal sensitivity following photorefractive keratectomy for myopia. *Journal of refractive surgery (Thorofare, NJ : 1995)*. 1999;15(1):38–45.
- [43] Linna TU, Pérez-Santonja JJ, Tervo KM, Sakla HF, Alió y Sanz JL, Tervo TM. Recovery of corneal nerve morphology following laser in situ keratomileusis. *Exp Eye Res*. 1998;66(6):755–763. doi:10.1006/exer.1998.0469.
- [44] Linna TU, Vesaluoma MH, Pérez-Santonja JJ, Petroll WM, Alió JL, Tervo TM. Effect of myopic LASIK on corneal sensitivity and morphology of subbasal nerves. *Invest Ophthalmol Vis Sci*. 2000;41(2):393–397.
- [45] Campos M, Hertzog L, Garbus JJ, McDonnell PJ. Corneal sensitivity after photorefractive keratectomy. *Am J Ophthalmol*. 1992;114(1):51–54.
- [46] Ishikawa T, Park SB, Cox C, del Cerro M, Aquavella JV. Corneal sensation following excimer laser photorefractive keratectomy in humans. *J Refract Corneal Surg*. 1994;10(4):417–422.
- [47] Richter A, Slowik C, Somodi S, Vick HP, Guthoff R. Corneal reinnervation following penetrating keratoplasty—correlation of esthesiometry and confocal microscopy. *Ger J Ophthalmol*. 1996;5(6):513–517.
- [48] Mathers WD, Jester JV, Lemp MA. Return of human corneal sensitivity after penetrating keratoplasty. *Arch Ophthalmol*. 1988;106(2):210–211.
- [49] Ruben M, Colebrook E. Keratoplasty sensitivity. *British Journal of Ophthalmology*. 1979;63(4):265–267.
- [50] Stamer L, Böhnke M, Draeger J. [Development of corneal sensitivity following keratoplasty]. *Fortschr Ophthalmol*. 1987;84(5):432–435.
- [51] Darwish T, Brahma A, Efron N, O'Donnell C. Subbasal nerve regeneration after penetrating keratoplasty. *Cornea*. 2007;26(8):935–940. doi:10.1097/ICO.0b013e3180de493f.
- [52] Al-Aqaba MA, Otri AM, Fares U, Miri A, Dua HS. Organization of the regenerated nerves in human corneal grafts. *Am J Ophthalmol*. 2012;153(1):29–37.e4. doi:10.1016/j.ajo.2011.06.006.
- [53] Draeger J, Schloot W, Wirt H. Interindividual differences of corneal sensitivity. Genetic aspects. *Ophthalmic paediatrics and genetics*. 1985;6(1-2):291–295.
- [54] Rosenberg ME, Tervo TM, Immonen IJ, Müller LJ, Gronhagen-Riska C, Vesaluoma MH. Corneal structure and sensitivity in type 1 diabetes mellitus. *Invest Ophthalmol Vis Sci*. 2000;41(10):2915–2921.
- [55] Tavakoli M, Petropoulos IN, Malik RA. Assessing corneal nerve structure and function in diabetic neuropathy. *Clin Exp Optom*. 2012;95(3):338–347. doi:10.1111/j.1444-0938.2012.00743.x.
- [56] Tavakoli M, Quattrini C, Abbott C, et al. Corneal confocal microscopy: a novel noninvasive test to diagnose and stratify the severity of human diabetic neuropathy. *Diabetes Care*. 2010;33(8):1792–1797. doi:10.2337/dc10-0253.
- [57] Tavakoli M, Kallinikos PA, Efron N, Boulton AJM, Malik RA. Corneal sensitivity is reduced and relates to the severity of neuropathy in patients with diabetes. *Diabetes Care*. 2007;30(7):1895–1897. doi:10.2337/dc07-0175.
- [58] Schwartz DE. Corneal sensitivity in diabetics. *Arch Ophthalmol*. 1974;91(3):174–178.
- [59] Rogell GD. Corneal hypesthesia and retinopathy in diabetes mellitus. *Ophthalmology*. 1980;87(3):229–233.
- [60] Murphy PJ. Noninvasive Assessment of Corneal Sensitivity in Young and Elderly Diabetic and Nondiabetic Subjects. *Invest Ophthalmol Vis Sci*. 2004;45(6):1737–1742. doi:10.1167/iovs.03-0689.
- [61] Tavakoli M, Mito-Pretorian M, Petropoulos IN, et al. Corneal confocal microscopy detects early nerve regeneration in diabetic neuropathy after simultaneous pancreas and kidney transplantation. *Diabetes*. 2013;62(1):254–260. doi:10.2337/db12-0574.
- [62] Paton L. The trigeminal and its ocular lesions. *The British Journal of Ophthalmology*. 1926:305–342.
- [63] Heigle TJ, Pflugfelder SC. Aqueous tear production in patients with neurotrophic keratitis. *Cornea*. 1996;15(2):135–138.
- [64] Wilson SE, Ambrósio R. Laser in situ keratomileusis-induced neurotrophic epitheliopathy. *Am J Ophthalmol*. 2001;132(3):405–406.

## Quellenverzeichnis für Tabelle und Abbildungen

- [1] Pflugfelder SC. Tear dysfunction and the cornea: LXVIII Edward Jackson Memorial Lecture. *Am J Ophthalmol*. 2011;152(6):900–909.e1. doi:10.1016/j.ajo.2011.08.023.
- [2] Pérez-Santonja JJ, Sakla HF, Cardona C, Chipont E, Alió JL. Corneal sensitivity after photorefractive keratectomy and laser in situ keratomileusis for low myopia. *Am J Ophthalmol*. 1999;127(5):497–504.
- [3] Murphy PJ, Corbett MC, O'Brart DP, Verma S, Patel S, Marshall J. Loss and recovery of corneal sensitivity following photorefractive keratectomy for myopia. *Journal of refractive surgery (Thorofare, NJ : 1995)*. 1999;15(1):38–45.
- [4] Linna TU, Vesaluoma MH, Pérez-Santonja JJ, Petroll WM, Alió JL, Tervo TM. Effect of myopic LASIK on corneal sensitivity and morphology of subbasal nerves. *Invest Ophthalmol Vis Sci*. 2000;41(2):393–397.
- [5] Benitez-del-Castillo JM, del Rio T, Iradier T, Hernández JL, Castillo A, Garcia-Sanchez J. Decrease in tear secretion and corneal sensitivity after laser in situ keratomileusis. *Cornea*. 2001;20(1):30–32.
- [6] Gallar J, Acosta MC, Moilanen JAO, Holopainen JM, Belmonte C, Tervo TMT. Recovery of corneal sensitivity to mechanical and chemical stimulation after laser in situ keratomileusis. *Journal of refractive surgery (Thorofare, NJ : 1995)*. 2004;20(3):229–235.
- [7] Bragheeth MA, Dua HS. Corneal sensation after myopic and hyperopic LASIK: clinical and confocal microscopic study. *British Journal of Ophthalmology*. 2005;89(5):580–585. doi:10.1136/bjo.2004.046888.
- [8] Darwish T, Brahma A, O'Donnell C, Efron N. Subbasal nerve fiber regeneration after LASIK and LASEK assessed by noncontact esthesiometry and in vivo confocal microscopy: prospective study. *J Cataract Refract Surg*. 2007;33(9):1515–1521. doi:10.1016/j.jcrs.2007.05.023.
- [9] Li M, Zhao J, Shen Y, et al. Comparison of Dry Eye and Corneal Sensitivity between Small Incision Lenticule Extraction and Femtosecond LASIK for Myopia. *PLoS ONE*. 2013;8(10):e77797. doi:10.1371/journal.pone.0077797.
- [10] Tavakoli M, Petropoulos IN, Malik RA. Assessing corneal nerve structure and function in diabetic neuropathy. *Clin Exp Optom*. 2012;95(3):338–347. doi:10.1111/j.1444-0938.2012.00743.x.