Weichlinsenmaterial – Angebot und Auswahl

Literatur

[1] Nichols JJ. Annual Report: Contact Lenses 2010. Contact Lens Spectrum 2011;Januar.

[2] Stapleton F, Keay L, Edwards K, et al. The Incidence of Contact Lens-Related Microbial Keratitis in Australia. Ophthalmology 2008;115:1655-1662.

[3] Dart JKG, Radford CF, Minassian D, et al. Risk Factors for Microbial Keratitis with Contemporary Contact Lenses. Ophthalmology 2008;115:1647-1654.

[4] Szczotka-Flynn, L., Ahmadian, R., & Diaz, M. (2009). A re-evaluation of the risk of microbial keratitis from overnight contact lens wear compared with other life risks Eye & contact lens, 35(2), 69–75. doi:10.1097/ICL.0b013e3181998dd3

[5] Morgan PB, Woods CA, Tranoudis IG, et al. International Contact Lens Prescribing in 2010. Contact Lens Spectrum 2010; January.

[6] Radford, CF, Minassian, D, Dart JFG, et al. Risk Factors for Nonulcerative Contact Lens Complications in an Ophthalmic Accident and Emergency Department. Ophthalmology 2009;116:385-392.

[7] Morgan P, Brennan, N. Evaluating Corneal Oxygenation during Lens Wear. Contact Lens Spectrum (Special Edition) 2007; May Special Edition.

[8] Morgan P, Brennan, N. Clinical highs and lows of Dk/t Part 2- Modulus, design, surface- more than just fresh air. Optician 2009; Juli.

[9] Chalmers RL, Wagner H, Mitchell GL, et al. Age and Other Risk Factors for Corneal Infiltrative and Inflammatory Events in Young Soft Contact Lens Wearers from the Contact Lens Assessment in Youth (CLAY) Study. Invest Ophthalmol Vis Sci 2011;52(9):6690-6.

[10] Brennan NA, Coles MLC, Comstock TL, Levy B. A 1-year prospective clinical trial of balafilcon A (PureVision) silicone-hydrogel contact lenses used on a 30-day continuous wear schedule. Ophthalmology. 2002;109:1172-1177.

[11] Chalmers RL, Hunt C, Hickson-Curran S, Young G. Struggle with hydrogel CL wear increases with age in young adults. Cont Lens Anterior Eye. 2009;32:113-119.

[12] Cheung SW, Cho P, Chan B, Choy C, Ng V. A comparative study of biweekly disposable contact lenses: Silicone hydrogel versus hydrogel. Clin Exp Optom. 2007;90:124-131.

[13] Dillehay SM, Miller MB. Performance of lotrafilcon B silicone hydrogel contact lenses in experienced low-Dk/t daily lens wearers. Eye Contact Lens. 2007;33:272-277.

[14] Dumbleton K, Keir N, Moezzi A, Feng Y, Jones L, Fonn D. Objective and subjective responses in patients refitted to daily-wear silicone hydrogel contact lenses. Optom Vis Sci. 2006;83:758-768.

[15] Guillon M, Maissa C. Use of silicone hydrogel material for daily wear. Cont Lens Anterior Eye. 2007;30:5-10.

[16] Riley C, Young G, Chalmers R. Prevalence of ocular surface symptoms, signs, and uncomfortable hours of wear in contact lens wearers: The effect of refitting with daily-wear silicone hydrogel lenses (senofilcon A). Eye Contact Lens. 2006;32:281-286.

[17] Santodomingo-Rubido J, Barrado-Navascues E, Rubido-Crespo M. Ocular surface comfort during the day assessed by instant reporting in different types of contact and non-contact lens wearers. Eye and Contact Lens. 2010;36:96-100.

[18] Schafer J, Mitchell GL, Chalmers RL, Long B, Dillehay S, Barr J, Bergenske P, Donshik P, Secor G, Yoakum J. The stability of dryness symptoms after refitting with silicone hydrogel contact lenses over 3 years. Eye Contact Lens. 2007;33:247-252.

[19] Young G, Riley CM, Chalmers RL, Hunt C. Hydrogel lens comfort in challenging environments and the effect of refitting with silicone hydrogel lenses. Optom Vis Sci. 2007;84:302-308.

[20] Papas E. Silicone Hydrogels: Is There a Comfort Advantage? Contact Lens Spectrum 2011; November.

[21] Young G, Riley CM, Chalmers RL, Hunt C. Hydrogel Lens Comfort in Challenging Environments and the Effect of Refitting with Silicone Hydrogel Lenses. Optom Vis Sci 2007;84:302-308.

[22] Richdale K, Sinnott LT, Skadahl E, Nichols JJ. Frequency of and Factors Associated With Contac Lens Dissatisfaction and Discontinuation. Cornea 2007;26:168-174.

1