

# Myopieprävention in der Praxis

Pascal Blaser, Nick Dash

## Literatur

- [1] N.S. Logan, P. Shah, A.R. Rudnicka, B. Gilmartin, C.G. Owen, Childhood ethnic differences in ametropia and ocular biometry: the Aston Eye Study, *Ophthalmic Physiol. Opt.* 31 (2011) 550–558.
- [2] Holden BA, Fricke TR, Wilson DA, Jong M, Naidoo KS, Sankaridurg P, Wong TY, Naduvilath TJ, Resnikoff S, Global Prevalence of Myopia and High Myopia and Temporal Trends from 2000 through 2050, *Ophthalmology*, May 2016 Volume 123, Issue 5, Pages 1036–1042.
- [3] Vitale, S., R.D. Sperduto, and F.L. Ferris, III, Increased Prevalence of Myopia in the United States Between 1971-1972 and 1999-2004. *Archives of Ophthalmology*, 2009. 127(12): p. 1632-1639.
- [4] Logan, N.S., et al., Childhood ethnic differences in ametropia and ocular biometry: the Aston Eye Study. *Ophthalmic Physiol Opt*, 2011. 31(5): p. 550-8.
- [5] O'Donoghue, L., et al., Refractive error and visual impairment in school children in Northern Ireland. *Br J Ophthalmol*, 2010. 94(9): p. 1155-9.
- [6] Katie M. Williams, et al, Prevalence of refractive error in Europe: the European Eye Epidemiology (E3) Consortium
- [7] McCullough, S. et al. „Six Year Refractive Change among White Children and Young Adults: Evidence for Significant Increase in Myopia among White UK Children.“ *PloS one* 11.1 (2016).
- [8] Vonghanit et al. Contact Lens & Anterior Eye BCLA Abstracts 2012, Blue Mountains Study.
- [9] Ryan SJ et al. *Retina*. 5th edition. Philadelphia: Elsevier; 2012:1115-1133.
- [10] Hayashi K et al. *Ophthalmology*. 2010; 117(8):1595-1611.
- [11] Flitcroft, D. I. „The complex interactions of retinal, optical and environmental factors in myopia aetiology.“ *Progress in retinal and eye research* 31.6 (2012): 622-660.
- [12] Virginie J. Verhoeven et al: Axial length and visual function in high myopia. ARVO poster 2016.
- [13] Parssinen, M. Kauppinen, A. Viljanen, The progression of myopia from its onset at age 8–12 to adulthood and the influence of heredity and external factors on myopic progression. A 23-year follow-up study, *Acta Ophthalmol.* 92 (2014) 730–739.
- [14] Blaser P. Myopia Care Available from: [www.myopiaprofile.org](http://www.myopiaprofile.org)
- [15] Gifford, K. Myopia Profile. 2015; Available from <http://www.myopiaprofile.com/> und <https://mykidsvision.org/>
- [16] Myopia Control. 2015; Available from: <http://www.myopiacontrol.org/>.
- [17] Richard L. Anderson, O.D. Myopia Prevention; Available from: <http://www.myopiaprevention.org/>
- [18] Rose KA, Morgan IG, Ip J, Kifley A, Huynh S, Smith W et al. Outdoor activity reduces the prevalence of myopia in children. *Ophthalmology*. 2009;115:1279–85.
- [19] Wu PC, Tsai CL, Wu HL, Yang YH, Kuo HK. Outdoor activity during class recess reduces myopia onset and progression in school children. *Ophthalmology*. 2013;120:1090–5.
- [20] J.C. Sherwin, M.H. Reacher, R.H. Keogh, A.P. Khawaja, D.A. Mackey, P.J. Foster, The association between time spent outdoors and myopia in children and adolescents a systematic review and meta-analysis, *Ophthalmology* 119 (2012) 2141–2151.
- [21] French, A.N. et al. „Time outdoors and the prevention of myopia.“ *Experimental eye research*. 114 (2013): 58-68.
- [22] D. Cheng, G.C. Woo, B. Drobe, K.L. Schmid, Effect of bifocal and prismatic bifocal spectacles on myopia progression in children three-year results of a randomized clinical trial, *JAMA Ophthalmol.* 132 (2014) 258–264.
- [23] J.J. Walline, K.L. Greiner, M.E. Mcvey, L.A. Jones-Jordan, Multifocal contact lens myopia control, *Optom. Vis. Sci.* 90 (2013) 1207–1214.
- [24] Aller TA, Liu M, Wildsoet CF, Myopia Control with Bifocal Contact Lenses: A Randomized Clinical Trial, *Optometry and vision science: official publication of the American Academy of Optometry* [Internet], 2016 Jan 18;93(4):1–352.
- [25] P. Cho, S.W. Cheung, Retardation of myopia in Orthokeratology (ROMIO) study: a 2-year randomized clinical trial, *Invest. Ophthalmol. Vis. Sci.* 53 (2012) 7077–7095.
- [26] Swarbrick H, The effects of overnight orthokeratology lens wear on corneal thickness. *Invest Ophthalmol Vis Sci.* 2003 Jun;44(6):2518-23.
- [27] J. Pauné, H. Morales, J. Armengol, L. Quevedo, M. Faria-Ribeiro, J.M. González- Méijome, Myopia control with a novel peripheral gradient soft lens and orthokeratology: a 2-year clinical trial, *Biomed. Res. Int.* 2015 (2015).
- [28] Turnbull PR, Munro OJ, Phillips JR. 'Contact Lens Methods for Clinical Myopia Control'. *Optom Vis Sci.* 2016 Sep; 93(9): 11206.
- [29] Huang, Jinhai, et al. „Efficacy comparison of 16 interventions for myopia control in children: A network meta-analysis.“ *Ophthalmology* 123.4 (2016): 697-709.
- [30] N.A. Brennan, Predicted reduction in high myopia for various degrees of myopia control, *Cont. Lens Anterior Eye* 35 (2012) e14–e15.
- [31] D. Adler, M. Millodot, The possible effect of undercorrection on myopic progression in children, *Clin. Exp. Optom.* 89 (2006) 315–321.
- [32] Wolf Lagrèze, Dr. med, Pressemitteilung anlässlich der DOG 2016
- [33] Chia, A., et al. „Atropine for the treatment of childhood myopia: safety and efficacy of 0.5%, 0.1%, and 0.01% doses (Atropine for the Treatment of Myopia 2)“ *Ophthalmology* 119.2 (2012): 347-354.
- [34] Chia A., et al. "Five-Year Clinical Trial on Atropine for the Treatment of Myopia 2: Myopia Control with Atropine 0.01% Eyedrops". *Ophthalmology*. 2016;123:391-9.
- [35] CooperVision; New contact lens therapy effective in slowing progression of juvenile-onset myopia, study shows; auf <http://www.news-medical.net/news/20161209/New-contact-lens-therapy-effective-in-slowng-progression-of-juvenile-onset-myopia-study-shows.aspx>