

Optometrische Möglichkeiten bei Diplopie nach Trauma

Vom Hirntumor zum Ironman

LITERATUR

- [1] Thiagarajan, P., Ciuffreda, K. J. (2014). Effect of Oculomotor Rehabilitation on Accommodative Responsivity in Mild Traumatic Brain Injury. *J. Rehabil. Res. Dev.*, 51, 175–192.
- [2] Ciuffreda, K. J., Ludlam, D. P., Yadav, N. K., Thiagarajan, P. (2016). Traumatic Brain Injury. *Adv. Ophthalmol. Optom.*, 1, 307–333.
- [3] Barton, J. J. S., Ranalli, P. J. (2020). Vision Therapy: Ocular Motor Training in Mild Traumatic Brain Injury. *Ann. Neurol.*, 88, 453–461.
- [4] Master, C. L., Scheiman, M., Gallaway, M., Goodman, A., Robinson, R. L., Master, S. R., Grady, M. F. (2016). Vision Diagnoses Are Common After Concussion in Adolescents. *Clin. Pediatr. (Phila.)*, 55, 260–267.
- [5] Green, W., Ciuffreda, K. J., Thiagarajan, P., Szymanowicz, D., Ludlam, D. P., Kapoor, N. (2010). Accommodation in Mild Traumatic Brain Injury. *J. Rehabil. Res. Dev.*, 47, 183.a
- [6] Capó-Aponte, J. E., Urosevich, T. G., Temme, L. A., Tarbett, A. K., Sanghera, N. K. (2012). Visual Dysfunctions and Symptoms During the Subacute Stage of Blast-Induced Mild Traumatic Brain Injury. *Mil. Med.*, 177, 804–813.
- [7] Scheiman, M., Wick, B. (2015). *Clinical Management of Binocular Vision*; Wolters Kluwer: Philadelphia, 2015.
- [8] Ciuffreda, K. J., Rutner, D., Kapoor, N., Suchoff, I. B., Craig, S., Han, M. E. (2008). Vision Therapy for Oculomotor Dysfunctions in Acquired Brain Injury: A Retrospective Analysis. *Optom. - J. Am. Optom. Assoc.*, 79, 18–22.
- [9] Kapoor, N., Ciuffreda, K. J. (2002). Vision Disturbances Following Traumatic Brain Injury. *Curr. Treat. Options Neurol.*, 4, 271–280.
- [10] Smaakjær, P., Wachner, L. G., Rasmussen, R. S. (2022). Vision Therapy Improves Binocular Visual Dysfunction in Patients with Mild Traumatic Brain Injury. *Neurol. Res.*, 44, 439–445.
- [11] Hunt, A. W., Mah, K., Reed, N., Engel, L., Keightley, M. (2016). Oculomotor-Based Vision Assessment in Mild Traumatic Brain Injury: A Systematic Review. *J. Head Trauma Rehabil.*, 31, 252–261.
- [12] Tamhankar, M. A., Ying, G., Volpe, N. J. (2012). Effectiveness of Prisms in the Management of Diplopia in Patients Due to Diverse Etiologies. *J. Pediatr. Ophthalmol. Strabismus*, 49, 222–228.
- [13] Santo, A. L., Race, M. L., Teel, E. F. (2020). Near Point of Convergence Deficits and Treatment Following Concussion: A Systematic Review. *J. Sport Rehabil.*, 29, 1179–1193.
- [14] Schroeder, T. L., Rainey, B. B., Goss, D. A., Grosvenor, T. P. (1996). Reliability of and Comparisons Among Methods of Measuring Dissociated Phoria. *Optom. Vis. Sci.*, 73, 389–397.
- [15] Tamhankar, M. A., Ying, G., Volpe, N. J. (2011). Success of Prisms in the Management of Diplopia Due to Fourth Nerve Palsy. *J. Neuroophthalmol.*, 31, 206–209.
- [16] Tamhankar, M. A., Ying, G.-S., Volpe, N. J. (2012). Prisms Are Effective in Re-solving Diplopia from Incomitant, Large, and Combined Strabismus. *Eur. J. Ophthalmol.*, 22, 890–897.
- [17] Anilkumar, S. E., Narendran, K. (2022). Prisms in the Treatment of Diplopia with Strabismus of Various Etiologies. *Indian J. Ophthalmol.*, 70, 609–612.
- [18] Gunton, K. B., Brown, A. (2012). Prism Use in Adult Diplopia. *Curr. Opin. Ophthalmol.*, 23, 400–404.
- [19] Hatt, S. R., Leske, D. A., Liebermann, L., Holmes, J. M. (2014). Successful Treatment of Diplopia with Prism Improves Health-Related Quality of Life. *Am. J. Ophthalmol.*, 157, 1209–1213.
- [20] Portela-Camino, J. A., Sanchez, I., Gutierrez, C., Martín-González, S. (2024). Successful Treatment of Diplopia Using Prism Correction Combined with Vision Therapy/Orthoptics Improves Health-Related Quality of Life. *PeerJ*, 12, e17315.
- [21] Conrad, J. S., Mitchell, G. L., Kulp, M. T. (2017). Vision Therapy for Binocular Dysfunction Post Brain Injury. *Optom. Vis. Sci. Off. Publ. Am. Acad. Optom.*, 94, 101–107.
- [22] Brautaset, R. L., Jennings, J. A. M. (2001). Associated Phoria and the Measuring and Correcting Methodology after H.-J. Haase (MKH). *Strabismus*, 9, 165–176.
- [23] Kushner, B. J. (2001). Recently Acquired Diplopia in Adults With Long-Standing Strabismus. *Arch. Ophthalmol.*, 119, 1795.
- [24] Lee, S. K., Zabrowski, C., Lee, M. S. (2025). Management of Diplopia Using Contact Lens. *Neuro-Ophthalmol.*, 49, 11–16.
- [25] Iacobucci, I. L., Furr, B. A., Archer, S. M. (2009). Management of Binocular Diplopia Due to Maculopathy with Combined Bangerter Filter and Fresnel Prism. *Am. Orthopt. J.*, 59, 93–97.
- [26] Rutstein, R. P. (2010). Use of Bangerter Filters with Adults Having Intractable Diplopia. *Optom. - J. Am. Optom. Assoc.*, 81, 387–393.
- [27] Newsham, D., O'Connor, A. R., Harrad, R. A. (2018). Incidence, Risk Factors and Management of Intractable Diplopia. *Br. J. Ophthalmol.*, 102, 393–397.
- [28] van Vliet, H., Jellema, H. M., Nieuwendaal, C., Lapid-Gortzak, R., Riemsdag, F., van der Meulen, I. (2021). Grey Filter Contact Lens as Therapeutic Option for Acquired Reduced Binocular Visual Performance. *Br. Ir. Orthopt. J.*, 17, 62–69.
- [29] Ciuffreda, K. J., Ludlam, D. P., & Kapoor, N. (2009). Clinical oculomotor training in traumatic brain injury. *Optometry and Vision Development*, 40(1), 16–23.